

THE BROAD Fe K LINE PROFILE IN NGC 4151

Jun-Xian Wang^{1,2}, You-Yuan Zhou^{1,2,3}, and Ting-Gui Wang^{1,2}

Received _____; accepted _____

¹Center for Astrophysics, University of Science and Technology of China, Hefei, Anhui, 230026, P. R. China; jxw@mail.ustc.edu.cn

²National Astronomical Observatories, Chinese Academy of Sciences

³Beijing Astrophysics Center, Beijing, 100080, P. R. China

ABSTRACT

We present an analysis of the Fe K line profile of NGC 4151 by using long ASCA observation data obtained in May 1995. The unprecedented good data quality, which is much better in the energy band around 6.4 keV than that of the famous 4.2-day ASCA observation of MCG –6-30-15 in July 1994, offers a unique opportunity to study the details of Fe K line profile. Apart from those characteristics already noticed in earlier ASCA observations on this object (Yaqoob et al. 1995): a broad and skewed profile, with a strong peak at about 6.4 keV and a large red wing extending to $\sim 4 - 5$ keV, which is remarkably similar to that of MCG –6-30-15, we also find a weak blue wing extending to about 8 keV, thanks to the good quality of the data. When fitted by a relativistic accretion disk line plus a narrow core at 6.4 keV, the data constrain the accretion disk to be nearly face-on, contrary to the edge-on geometry inferred from optical and UV observations. However, the extended blue wing can not be well fitted even after we include corresponding Fe K β components. Ni K α line emission by an amount of 12% of Fe K α is statistically required. An alternative explanation is a model consisting of a narrow core and two disk lines with inclinations of 58° and 0° , respectively. We suppose that the component with inclination of 58° was observed directly, consistent with its edge-on geometry, and the component with inclination of 0° was scattered into our line of sight by a Compton mirror, which might be the cool accretion disk corona proposed by Poutanen et al. (1996).

Subject headings: black hole physics — galaxies: active — galaxies: individual (NGC 4151) — line: profiles — X-rays: galaxies

1. Introduction

The fluorescence iron $K\alpha$ line revealed by a 4.2-day ASCA observation in the Seyfert 1 galaxy MCG –6-30-15 has distinct features which are unique characteristics of the emission line from a relativistic disk (Tanaka et al. 1995). The line is peaked at around 6.4 keV, with a broad red wing extending down to 4 keV. The skewness of the profile towards red and the sharp blue edge near the rest frame energy are a consequence of the combination of relativistic Doppler effect and strong gravitational redshift, and are unique to a low-inclination relativistic accretion disk; other mechanisms of the line formation are less successful to produce these features (Fabian et al. 1995). However, the alternative models such as thermal Comptonization (Misra and Kembhavi 1998) can not be ruled out completely. Both the disk line fit (Tanaka et al. 1995) and the frequency extrema analysis (Bromley, Miller & Pariev 1998) suggested that the inner edge of line formation region is within a few Schwarzschild radii, providing ever strongest evidence for a super-massive black hole in this object. Similar broad line profiles with lower statistics have also been seen in other AGNs (Mushotzky et al. 1995, Tanaka et al. 1995, Yaqoob et al. 1995, Nandra et al. 1997, Turner et al. 1998 and references therein).

The line profile is strongly dependent on the inclination of the relativistic accretion disk (Fabian et al. 1989), thus providing a method to estimate the inclination of inner accretion disk from line profile fitting. The inclinations from this method for a sample of Seyfert 1 galaxies are strongly constrained to be nearly face-on (Nandra et al. 1997), consistent with the expectation of the unification model of the two types of Seyfert galaxies (see Antonucci 1993 for a review). However, there are complications to this simple picture. Turner et al. (1998) found that type 2 Seyfert galaxies also possess similar iron $K\alpha$ line profiles, indicative of face-on accretion disk as well. This might be due to the strong contribution of a narrow component in these objects (Weaver & Reynolds 1998) or due to complex

absorption (Wang et al. 1999a). Moreover, the rapid variability of the line equivalent width (EW) and profile observed in MCG –6-30-15 (Iwasawa et al. 1996) and NGC 4051 (Wang et al. 1999b) cannot be readily explained by any current simple accretion disk line model.

NGC 4151 is a bright nearby ($z = 0.0033$) Seyfert 1.5 galaxy. The edge-on orientation of its nucleus is strongly supported by the biconical geometry of the [O III] $\lambda 5007$ region (Evans et al. 1993, Pedlar et al. 1993) and the best estimated inclination is 65° . Yaqoob et al. (1995) presented the first measurement of its broad Fe K line profile from the ASCA observations performed in 1993 May, November and December. The apparent line profile is asymmetric, consisting of a peak at $\sim 6.3 - 6.4$ keV and a broad red wing extending to $\sim 4 - 5$ keV. When a disk line model is fitted, the inclination angle of the disk ($\theta = 0_{-0}^{+19}$ degrees) is strongly constrained to be face-on, in contrary to the edge-on geometry of this source. This problem can be eased, but not solved, by assuming an additional narrow-line contribution, presumably from the torus, to the Fe K line ($\theta = 25_{-8}^{+3}$ degrees, see Yaqoob & Weaver 1997).

A much longer ASCA observation (3 days) of NGC 4151 was carried out in 1995, which provides data with better statistics (Leightly et al. 1997). In this letter, we report measurement of the complex broad iron K line profile and its implication.

2. The ASCA Data

NGC4151 was observed by ASCA from 1995 May 10 to 12 with the Solid-state Imaging Spectrometer (SIS) in 1CCD mode and the Gas Imaging Spectrometer (GIS) in PH mode. The GIS data are contaminated by a nearby BL Lac object due to their worse spatial resolution. In this letter, we will concentrate on the SIS data, also because they offer much better energy resolution (Inoue 1993), which is crucial to the Fe K line profile analysis.

The data were reduced with the ASCA standard software XSELECT and cleaned using the following screening criteria: satellite not passing through the South Atlantic Anomaly (SAA), geomagnetic cutoff rigidity greater than 6 GeVc^{-1} , and minimum elevation angle above Earth’s limb of 10° and 20° for nighttime and daytime observations, respectively. Source counts were extracted from a circular area of radius $3.4'$ for the SIS0 and SIS1, and the background counts were estimated from the blank sky data. Spectra extracted from SIS0 and SIS1 were combined together and grouped to have >20 counts in each bin to allow the use of χ^2 statistics. Spectral analysis was carried out using XSPEC.

The ASCA observation lasted for about three days, and the SIS0 detector received a net exposure time of 93,000s and an average count rate of 1.49 cts/s in the $0.4 - 10.0 \text{ keV}$ band, while these two parameters for the well-known 4.2-day ASCA observation on MCG –6-30-15 in 1994 July are 150,000s and 1.82 cts/s, respectively. Although shorter exposure time and lower count rate may probably lead to worse statistics, we noticed that, because of its much harder spectrum (due to strong absorption), the actual total counts (32,000) for NGC 4151 in the $5.0 - 7.0 \text{ keV}$ band are almost three times of those (11,000) for MCG –6-30-15 in the same energy band. Thus, the NGC 4151 Fe K spectrum used in this paper has better statistics than the average MCG –6-30-15 spectrum and a best ever quality Fe K line profile.

3. Spectral fits

Following Weaver et al. (1994), we fit the underlying continuum in the $1.0 - 4.0$ and $8.0 - 10.0 \text{ keV}$ bands (to exclude the possible broad iron line region) with a model which consists of a dual absorbed power law with some fraction (a best-fit value for this letter is $\sim 5\%$) of the direct continuum scattered into our line of sight and absorbed only by the Galactic column of $2 \times 10^{20} \text{ cm}^{-2}$. We do not include a Compton reflection

component in the spectral fits because a clear reflection component has never been detected (Maisack & Yaqoob 1991; Yaqoob et al. 1993). Zdziarski et al. (1996) claimed that they detected a reflection component. However, considering the poor statistical significance of their detection (they used only a delta-chi square of 2.7 for the their errors yet the model was complicated), the complex intrinsic continuum which was unknown, and the cross-calibration of two different satellites (Ginga and GRO/OSSE) which is not perfect, we think that their result is questionable.

Our model can describe the data reasonably well ($\chi^2 = 364$ for 325 degrees of freedom), and the best-fit model parameters converge to $\Gamma = 1.30 \pm 0.07$, $N_H(\text{low}) \simeq 3.2 \times 10^{22} \text{ cm}^{-2}$ (covering $\sim 40\%$ of the source), and $N_H(\text{high}) = \sim 10.9 \times 10^{22} \text{ cm}^{-2}$ (covering $\sim 60\%$ of the source). The index is slightly flatter than that seen previously for NGC 4151 as derived from data weighted toward higher energies (Yaqoob et al. 1993). We have also tried a model consisting of a power-law absorbed by ionized material (Zdziarski et al. 1995) plus a fraction of scattered underlying continuum, but failed to get a satisfactory fit ($\Gamma = 1.00$, $\chi^2 = 393$ for 326 degrees of freedom).

The profile of the iron K line is shown in Figure 1a, which is similar to, and much better defined than, the one presented by Yaqoob et al. (1995). The line shows a strong narrow peak around 6.4 keV and a huge red wing, containing a second slightly weak peak at around 5.2 keV, extending to ~ 4.5 keV. In this respect, the line profile of NGC 4151 is remarkably similar to that seen in MCG –6-30-15 (see Figure 1b for comparison). We also find a weak blue wing extending to about 8 keV, which is just visible in Figure 2c of Yaqoob et al. (1995), is now clearly seen, thanks to the greatly improved statistics. The brightness of this source rules out the possibility that the blue wing is caused by improper background subtraction (only 1% of the count rate at 7. – 8. keV is due to the background).

First, we fit the line with a disk-line model (Fabian et al. 1989) plus a narrow core at

6.4 keV which is presumably from the torus and also statistically required ($\Delta\chi^2 = -46$). Though expected in theory, the fluorescence Fe K β line is always ignored when fitted to data with much lower quality. Considering the high quality of the data in this letter, we also include the corresponding Fe K β components (disk-line plus narrow core) in all of our models by amounts of 11.3% of Fe K α (George & Fabian 1991). In fact, Fe K β components are also statistically required here ($\Delta\chi^2 = -36$). The outer radius (R_o) of the disk is fixed at $1000 R_g$ ($R_g = GM/c^2$) to minimize the numbers of free parameters because the fits are not sensitive to the value for the outer radius. The best fit value is also consistent with $1000 R_g$ when it is allowed to vary. Considering the strong peak at 6.4 keV, we fixed the disk-line energy at 6.4 keV in the source rest frame. Results of this fit are given in Table 1 (Model A). The inclination of the disk is 27° , similar to the value obtained by Yaqoob & Weaver (1997). However, the blue-wing of the line beyond the 6.8 keV has not been well fitted (see Figure 2a).

In the accretion disk model context, the high energy blue wing can be produced by an accretion disk with high inclination angle. The blue wing, as well as the edge-on geometry of NGC 4151 inferred from optical and UV observations (Evans et al. 1993), motivated us to add an extra component of a disk line with large inclination to Model A (Model B, Fe K β components also included). We assume the inner and outer radii of the disk for these two disk-line components are the same, while the emissivity index q ($F_{line} \propto R^{-q}$) is allowed to vary independently considering the fact the line photons from different parts of the disk may have different scattering fraction. The results are also given in Table 1. The model can fit the observed line profile fairly well (see Figure 2b). The improvement of Model B to Model A is significant ($\Delta\chi^2 = -19$). As expected, the second disk component requires a high inclination of 58_{-12}^{+32} degrees, while the first component requires a face-on disk (0_{-0}^{+12} degrees).

Alternatively, the high energy blue wing between 6.8 and 8.0 keV may also be produced by the Ni K α line emission at 7.48 keV. In order to reduce the excesses in the residuals between 6.8 and 8.0 keV in Figure 2a, corresponding Ni K α components by amounts of $12^{+6}_{-6}\%$ of Fe K α have to be added to Model A (see Figure 2c, $\Delta\chi^2 = -13$).

4. Discussion

Zdziarski et al. (1996) argued that the ASCA data of NGC 4151 obtained in May 1993 can be modeled by complex absorption and a narrow Fe K line only, with no broad line required, contrary to the results of Yaqoob et al. (1995) and this letter. However, the ASCA data used by Zdziarski et al have much lower signal-to-noise. Through fitting the unprecedented high quality spectra of NGC 4151 obtained by ASCA in 1995, we find a prominent broad Fe K line profile in NGC 4151 (see Figure 1), which is remarkably similar to that of MCG –6-30-15. A model consisting a dual absorbed power law plus a scattered component and a narrow Fe K line at 6.4 keV ($\sigma = 0.11^{+0.02}_{-0.02}$) fits the data (1.0 – 10.0 keV) poorly ($\chi^2 = 706$ for 598 degrees of freedom) and results broad systematic positive residuals around 6.0 keV (see Figure 2d). An extra broad line is statistically required ($\Delta\chi^2 = -78$).

The high signal-to-noise profile of NGC 4151 obtained in this letter shows a strong narrow peak around 6.4 keV and a huge red wing, containing a second slightly weak peak at around 5.2 keV, extending to ~ 4.5 keV. In this respect, the line profile of NGC 4151 is remarkably similar to that seen in MCG –6-30-15. We also find a weak blue wing extending to about 8.0 keV, which is just visible in Figure 2c of Yaqoob et al. (1995), is now clearly seen (Figure 1a), thanks to the greatly improved statistics. There may also be such weak blue wing in the Fe K profile of MCG –6-30-15 (see Figure 1b).

When fitted by the disk-line model plus a narrow core, a face-on disk is also required (θ

$= 27^\circ$), contrary to the edge-on geometry inferred from optical and UV observations (Evans et al. 1993, Pedlar et al. 1993). We consider that the Compton mirror of Poutanen et al. (1996), who proposed that the central source of NGC 4151 is completely hidden from our line of sight and observed X-ray radiation is produced by scattering in the higher, cooler parts of the accretion disk corona, or in a wind, could solve the problem well. We also want to point out that scattering by an "ionizing cone" would also explain the similarity of Fe K line profiles seen in Seyfert 2 galaxies and in Seyfert 1 galaxies (Turner et al. 1998), indicating reprocessing by face-on disks and contradicting the expectation of the unification scheme. The variability properties of the Fe K line are important to test such scattering hypothesis.

Besides the broad profile of the Fe K line, the blue wing extending to about 8.0 keV (see Figure 1a), which could not be well fitted by a simple disk line plus a narrow core even after the corresponding Fe $k\beta$ components included, should also be paid much attention to. One possible explanation is that we observe not only a broad Fe K line in NGC 4151, but also a corresponding Ni $K\alpha$ line by an amount of $12^{+6}_{-6}\%$ of Fe $K\alpha$. This may be the first detection of Ni $K\alpha$ fluorescence line emission in the X-ray spectrum of AGNs.

However, we notice that the energy of Ni $K\alpha$ line is just above that of the Fe K absorption edge. The line flux, which is sensitive to the Ni abundance and X-ray spectral slope, can be substantially reduced due to photo-electron absorption by Fe atoms. So, the contribution of Ni $K\alpha$ emission line might be much weaker than expected. An alternative explanation to the weak blue wing is an extra Fe K disk line with high inclination angle (it is also possible that both the Ni $K\alpha$ components and the extra disk line are important in reality). Poutanen et al. (1996) assumed that the central source is completely hidden from our line of sight by the thick part of the accretion disk wind or by the broad emission-line clouds. However, it is plausible to suppose that the thick material covers the X-ray only

partially, and part of the central continuum is seen directly. The fast variability and the complex absorption of the X-ray continuum seem to support this idea. The results of two-disk component model fit (Model B) is fully consistent with this picture. Poutanen et al. (1996) assumed that the scattering region is situated along the axis of the accretion disk in the form of a cone. The half opening angle of the "ionized cone" determined from the optical and UV observation is about 35 degree (see Fig. 10 of Evans et al. 1993), the angle-averaged line profile in the scattered light should resemble the one from a face-on disk, which corresponds to the low inclination line component in our two disk-line fit. The direct component should show the same inclination as the system, which is 65° inferred from optical and UV observations. We got an inclination of 58_{-12}^{+32} degrees for this line component, which is in good agreement with interpreting it as a direct component. Comparable EW of the two line components suggests that the flux of the X-ray scattered should be commensurate roughly with the directly observed flux. The little lower parameter q of the direct disk-line indicates that more direct Fe K photons can be observed from large disk radii.

This work is supported by Chinese National Natural Science Foundation, PanDeng Project and Foundation of Ministry of Education. We thank the referee for many critical comments, especially on the possibility of Ni K α contribution, which significantly improve the presentation of this paper.

REFERENCES

- Antonucci, R. 1993, ARA&A, 31, 473
- Bromley, B. C., Miller, W. A. & Pariev, V. I. 1998, Nature, 391, 54
- Evans, I. N., et al., 1993, ApJ, 417, 82
- Fabian, A. C., Rees, M. J., Stella, L. & White, N. E. 1989, MNRAS, 238, 729
- Fabian, A. C., et al. 1995, MNRAS, 277, L11
- George, I.M. & Fabian, A.C. 1991, MNRAS, 249, 352
- Inoue, H. 1993, Exp.Astro., 4, 1
- Iwasawa, K., et al. 1996, MNRAS, 282, 1038
- Leighly, K. M., et al. 1997, X-ray Imaging and spectroscopy of cosmic hot Plasmas, Univ. Acad. Press, 291
- Maiscak, M. & Yaqoob, T. 1991, A&A, 249, 25
- Misra, R. & Kembhavi, A. K. 1998, ApJ, 499, 205
- Mushotzky, R. F., et al. 1995, MNRAS, 272, L9
- Nandra, K., George, I. M., Mushotzky, R. F., Turner, T.J. & Yaqoob, T. 1997, ApJ, 477, 602
- Pedlar, A., et al. 1993, MNRAS, 263, 471
- Poutanen, J., Sikora, M., Begelman, M. C. & Magdziarz, P. 1996, ApJ, 465, L107
- Tanaka, Y., et al. 1995, Nature, 375, 659
- Turner, T. J., George, I. M., Nandra, K. & Mushotzky, R. F. 1998, ApJ, 493, 91
- Wang, T.G., Mihara, T., Otani, C., Matsuoka, M., & Awaki, H., 1999a, ApJ in press
- Wang, J. X., Zhou, Y. Y., Xu, H. G. & Wang, T. G. 1999b, ApJ, 516, L65

- Weaver, K. A., et al. 1994, ApJ, 436, L27
- Weaver, K. A. & Reynolds, C. S. 1998, ApJ, 503, L39
- Yaqoob, T. & Weaver, K. A. 1997, X-ray Imaging and Spectroscopy of Cosmic Hot Plasmas, Univ. Acad. Press, 201
- Yaqoob, T., et al. 1993, MNRAS, 262, 435
- Yaqoob, T., et al. 1995, ApJ, 453, L81
- Zdziarski, A., et al. 1995, ApJ, 363, L1
- Zdziarski, A., et al. 1996, MNRAS, 283, 193




Fig. 1.— (a) The high S/N ratio Fe K line profile in NGC 4151. (b) The profile of Fe K line in MCG -6-30-15 (from SIS data of the well-known 4.2 days ASCA observation in 1994). It's obvious that the profile of Fe K line in NGC 4151 has a much higher S/N ratio.




Fig. 2.— Plots of residuals for models: (a) a single cold disk-line plus a narrow core (Model A); (b) two cold Fe K disk-line with different inclinations plus a narrow core (Model B); (c) including corresponding Ni $K\alpha$ components by amounts of 12% of Fe $K\alpha$ to Model A. Fe $K\beta$ components by amounts of 11.3% of Fe $K\alpha$ are included in both Model A and Model B; (d) a model consisting a dual absorbed power law plus a scattered component and a narrow Fe K line at 6.4 keV.

Table 1: Fe K line fits^a

Model	χ^2/dof	R_i (R_g)	EW_{core} (eV)	EW_{disk} (eV)	q	θ
A	628/599	$8.9^{+0.9}_{-0.9}$	71^{+13}_{-12}	276^{+21}_{-24}	$-4.2^{+0.7}_{-0.8}$	27^{+1}_{-2}
B	609/596	$6.0^{+0.8}_{-0.0}$	89^{+11}_{-10}	136^{+28}_{-27}	$-2.0^{+0.4}_{-0.3}$	58^{+32}_{-12}
				150^{+18}_{-19}	$-3.3^{+0.3}_{-0.2}$	0^{+12}_{-0}

^aErrors are given at 90% level. The energy of narrow core is fixed at 6.4 keV in the galaxy rest frame and the line width fixed at zero. Model A: a single cold disk line (6.4 keV in the galaxy rest frame) plus a narrow core; Model B: two cold disk line with different inclinations plus a narrow core. Fe K β components by amounts of 11.3% of Fe K α are included in both models.